

Interactions Between Asphaltenes and a Model Demulsifier in Bulk and at an Interface Studied by Small-angle Neutron Scattering (SANS) and Neutron Reflectometry

Kenneth D. Knudsen,^{†,||} Sébastien Simon,^{*,‡} Thomas Geue[#], Joshaniel F. K. Cooper[‡], Johan Sjöblom[⊥]

[†]Institute for Energy Technology (IFE), N-2007 Kjeller, Norway

^{||}Department of Physics, the Norwegian University of Science and Technology (NTNU), N-7491 Trondheim, Norway

[⊥]Ugelstad Laboratory, Department of Chemical Engineering, the Norwegian University of Science and Technology (NTNU), N-7491 Trondheim, Norway

[#] Laboratory for Neutron Scattering and Imaging (LNS), Paul Scherrer Institute, CH-5232 Villigen PSI, Switzerland

[‡] ISIS Neutron and Muon Source, Science and Technology Facilities Council, Rutherford Appleton Laboratory, Harwell Science and Innovation Campus, Oxon OX11 0QX, United Kingdom




Figure S1. SANS data for asphaltenes at 2.5 g/L in D-toluene. The data for 2.5 g/L asphaltenes was not model fitted, but the overall intensity is seen to scale quite well with the reduction in concentration compared to the 10 g/L (shown in the main text).